

Operations on polynomials

$p, q \in R[x], \deg(p) = \deg(q) = n$

- **Coefficient representation**

Addition: $O(n)$

Multiplication: $O(n^2) \longrightarrow O(n \log n)$

Evaluation at x_0 : $O(n)$

- **Point-value representation**

$$p = (x_0, y_0), (x_1, y_1), \dots, (x_n, y_n)$$

$$q = (x_0, z_0), (x_1, z_1), \dots, (x_n, z_n)$$

Operations on polynomials

Addition:

$$p + q = (x_0, \underbrace{y_0 + z_0}_{}), (x_1, \underbrace{y_1 + z_1}_{}), \dots, (x_n, \underbrace{y_n + z_n}_{})$$

Running time: $O(n)$

Multiplication:

$$p \cdot q = (x_0, \underbrace{y_0 \cdot z_0}_{}), (x_1, \underbrace{y_1 \cdot z_1}_{}), \dots, (x_n, \underbrace{y_n \cdot z_n}_{})$$

(Condition: $n \geq \text{degree}(pq)$)

Running time: $O(n)$

Evaluation at point x' : ??

Convert polynomial to coefficient representation
(interpolation)

Polynomial multiplication

Compute the product of two polynomials p, q of degree $< n$:

p, q of degree $n-1$, n coefficients



Evaluation: $x_0, x_1, \dots, x_{2n-1}$

$\mathcal{O}(n \log n)$
FFT

$2n$ point-value pairs $(x_i, p(x_i))$ und $(x_i, q(x_i))$



Pointwise multiplication

$\mathcal{O}(n)$

$2n$ point-value pairs $(x_i, pq(x_i))$



Interpolation

?

pq of degree $2n-2$, $2n-1$ coefficients

Divide-and-conquer approach

Idea: (assume n is even)

$$\begin{aligned}
 p(x) &= a_0 + a_1x + \dots + a_{n-1}x^{n-1} \\
 &\stackrel{\text{even}}{\longrightarrow} = a_0 + a_2x^2 + \dots + a_{n-2}x^{n-2} + \\
 &\stackrel{\text{odd}}{\longrightarrow} a_1x + a_3x^3 + \dots + a_{n-1}x^{n-1} \\
 &= a_0 + a_2x^2 + \dots + a_{n-2}(x^2)^{(n-2)/2} + \\
 &\quad x(a_1 + a_3x^2 + \dots + a_{n-1}(x^2)^{(n-2)/2}) \\
 &= p_0(x^2) + xp_1(x^2)
 \end{aligned}$$

$$p_0(x) = a_0 + a_2x + \dots + a_{n-2}x^{(n-2)/2}$$

$$p_1(x) = a_1 + a_3x + \dots + a_{n-1}x^{(n-2)/2}$$

Select x_0, \dots, x_{2n-1} such that the computations of $p(x_k)$ and $p(x_{k+n})$ are almost identical.

Representation of $p(x)$

Assume: $\text{degree}(p) < n$


3a. Values of the n powers of the principal n th root of unity

$$\omega_n = e^{2\pi i / n}$$

$$i = \sqrt{-1} \quad e^{2\pi i} = 1$$

Powers of ω_n (roots of unity):

$$1 = \underbrace{\omega_n^0, \omega_n^1, \dots, \omega_n^{n-1}}$$



Discrete Fourier Transform

The values $p(\omega_n^i)$ of the n powers of ω_n uniquely define p if $\text{degree}(p) < n$.

Discrete Fourier Transform (DFT)

$$DFT_n(p) = (p(\omega_n^0), p(\omega_n^1), \dots, p(\omega_n^{n-1}))$$

Example: $n = 4$

$$e^{ix} = \cos x + i \sin x$$


$$\omega_4^0 = e^{0i} = \cos(0) + i \sin(0) = 1$$

$$\omega_4^1 = e^{2\pi i / 4} = \cos(\pi / 2) + i \sin(\pi / 2) = i$$

$$\omega_4^2 = (e^{2\pi i / 4})^2 = \cos \pi + i \sin \pi = -1$$

$$\omega_4^3 = (e^{2\pi i / 4})^3 = \cos(3\pi / 2) + i \sin(3\pi / 2) = -i$$

Evaluation at the roots of unity



Evaluation at the roots of unity

$$p(x) = 3x^3 - 15x^2 + 18x$$

$$(\omega_4^0, p(\omega_4^0)) = (1, p(1)) = (1, 6)$$

$$(\omega_4^1, p(\omega_4^1)) = (i, p(i)) = (i, 15 + 15i)$$

$$(\omega_4^2, p(\omega_4^2)) = (-1, p(-1)) = (-1, -36)$$

$$(\omega_4^3, p(\omega_4^3)) = (-i, p(-i)) = (-i, 15 - 15i)$$

$$DFT_4(p) = (6, 15+15i, -36, 15-15i)$$

Polynomial multiplication

Compute the product of two polynomials p, q of degree $< n$:

p, q of degree $n-1$, n coefficients



Evaluation: $\underbrace{\omega_{2n}^0, \omega_{2n}^1, \dots, \omega_{2n}^{2n-1}}$

$2n$ point-value pairs $(\omega_{2n}^i, p(\omega_{2n}^i))$ and $(\omega_{2n}^i, q(\omega_{2n}^i))$



Pointwise multiplication

$2n$ point-value pairs $(\omega_{2n}^i, pq(\omega_{2n}^i))$



Interpolation

pq of degree $2n-2$, $2n-1$ coefficients

4. Properties of the roots of unity

$\omega_{2n}^0, \omega_{2n}^1, \dots, \omega_{2n}^{2n-1}$ form a **multiplicative group**

Cancellation lemma:

For any integers $n > 0$, $k \geq 0$ and $d > 0$ we have:

$$\underbrace{\omega_{dn}^{dk}}_{= \omega_n^k} = \omega_n^k$$

Proof:

$$\omega_{dn}^{dk} = e^{2\pi i dk / (dn)} = e^{2\pi i k / n} = \omega_n^k$$

Therefore:

$$\omega_{2n}^n = \omega_2^1 = -1$$

Group

$(G, *)$ G set, $*$ Operation

(1) Closure

(2) Associativity

(3) Neutral Element

(4) Inverse Element

Group of roots of unity

$G = \{ \omega_{2n}^0, \omega_{2n}^1, \dots, \omega_{2n}^{2n-1} \}$ * Complex multiplication

(1) Closure

(2) Associativity

(3) Neutral Element

(4) Inverse Element

5. Discrete Fourier Transform

$$DFT_n(p) = (p(\omega_n^0), p(\omega_n^1), \dots, p(\omega_n^{n-1}))$$

Fast Fourier Transform:

Computation of $DFT_n(p)$ by means of a divide-and-conquer approach.

Discrete Fourier Transform

Idea: (assume n is even)

$$\begin{aligned}
 p(x) &= a_0 + a_1x + \dots + a_{n-1}x^{n-1} \\
 &= a_0 + a_2x^2 + \dots + a_{n-2}x^{n-2} + \\
 &\quad a_1x + a_3x^3 + \dots + a_{n-1}x^{n-1} \\
 &= a_0 + a_2x^2 + \dots + a_{n-2}(x^2)^{(n-2)/2} + \\
 &\quad x(a_1 + a_3x^2 + \dots + a_{n-1}(x^2)^{(n-2)/2}) \\
 &= p_0(x^2) + xp_1(x^2)
 \end{aligned}$$

$$p_0(x) = a_0 + a_2x + \dots + a_{n-2}x^{(n-2)/2}$$

$$p_1(x) = a_1 + a_3x + \dots + a_{n-1}x^{(n-2)/2}$$

Discrete Fourier Transform

Putting things together:

$$DFT_n(p) = (p(\omega_n^0), p(\omega_n^1), \dots, p(\omega_n^{n-1}))$$

$$p(x) = p_0(x^2) + x \cdot p_1(x^2)$$

$$x = \omega_n^k, \quad (\omega_n^k)^2 = \omega_{n/2}^k \quad k = 0, \dots, n-1$$

$$\begin{aligned} (DFT_n(p))_k &= p(\omega_n^k) \\ &= p_0((\omega_n^k)^2) + \omega_n^k \cdot p_1((\omega_n^k)^2) \\ &= p_0(\omega_{n/2}^k) + \omega_n^k \cdot p_1(\omega_{n/2}^k) \\ &= \begin{cases} p_0(\omega_{n/2}^k) + \omega_n^k \cdot p_1(\omega_{n/2}^k) & k < \frac{n}{2} \\ p_0(\omega_{n/2}^{k-\frac{n}{2}}) + \omega_n^k \cdot p_1(\omega_{n/2}^{k-\frac{n}{2}}) & k \geq \frac{n}{2} \end{cases} \end{aligned}$$

$$\begin{aligned} &= \begin{cases} (DFT_{\frac{n}{2}}(p_0))_k + \omega_n^k (DFT_{\frac{n}{2}}(p_1))_k & k < \frac{n}{2} \\ (DFT_{\frac{n}{2}}(p_0))_{k-\frac{n}{2}} + \omega_n^k (DFT_{\frac{n}{2}}(p_1))_{k-\frac{n}{2}} & k \geq \frac{n}{2} \end{cases} \end{aligned}$$

Discrete Fourier Transform

Evaluation for $k = 0, \dots, n - 1$:

$$p(\omega_n^k) = p_0((\omega_n^k)^2) + \omega_n^k p_1((\omega_n^k)^2) = \begin{cases} p_0(\omega_{n/2}^k) + \omega_n^k p_1(\omega_{n/2}^k), & \text{if } k < n/2 \\ p_0(\omega_{n/2}^{k-n/2}) + \omega_n^k p_1(\omega_{n/2}^{k-n/2}), & \text{if } k \geq n/2 \end{cases}$$

$$\begin{aligned} DFT_n(p) &= (p_0(\omega_{n/2}^0), \dots, p_0(\omega_{n/2}^{n/2-1}), \quad p_0(\omega_{n/2}^0), \dots, p_0(\omega_{n/2}^{n/2-1})) \\ &\quad + (\omega_n^0 p_1(\omega_{n/2}^0), \dots, \omega_n^{n/2-1} p_1(\omega_{n/2}^{n/2-1}), \quad \omega_n^{n/2} p_1(\omega_{n/2}^0), \dots, \omega_n^{n-1} p_1(\omega_{n/2}^{n/2-1})) \end{aligned}$$

Discrete Fourier Transform

Example:

$$p(\omega_4^0) = p_0(\omega_2^0) + \omega_4^0 p_1(\omega_2^0)$$

$$p(\omega_4^1) = p_0(\omega_2^1) + \omega_4^1 p_1(\omega_2^1)$$

$$p(\omega_4^2) = p_0(\omega_2^0) + \omega_4^2 p_1(\omega_2^0)$$

$$p(\omega_4^3) = p_0(\omega_2^1) + \omega_4^3 p_1(\omega_2^1)$$

Computation of DFT_n

$$DFT_n(p) = (p(\omega_n^0), p(\omega_n^1), \dots, p(\omega_n^{n-1}))$$

Base case: $n = 1$ ($\text{degree}(p) = n - 1 = 0$)

$$DFT_1(p) = a_0$$

General case :

Divide:

Divide p into p_0 and p_1

Conquer:

Recursively compute $DFT_{n/2}(p_0)$ and $DFT_{n/2}(p_1)$.

Merge:

For $k = 0, \dots, n - 1$ compute:

$$\begin{aligned} DFT_n(p)_k = & (DFT_{n/2}(p_0), DFT_{n/2}(p_0))_k + \\ & \omega_n^k \cdot (DFT_{n/2}(p_1), DFT_{n/2}(p_1))_k \end{aligned}$$

A further improvement

$$\begin{aligned}
 p(\omega_n^k) &= \begin{cases} p_0(\omega_{n/2}^k) + \omega_n^k p_1(\omega_{n/2}^k) & \text{if } k < n/2 \\ p_0(\omega_{n/2}^{k-n/2}) + \cancel{\omega_n^k} p_1(\omega_{n/2}^{k-n/2}) & \text{if } k \geq n/2 \end{cases} \\
 &= \begin{cases} p_0(\omega_{n/2}^k) + \omega_n^k p_1(\omega_{n/2}^k) & \text{if } k < n/2 \\ p_0(\omega_{n/2}^{k-n/2}) - \cancel{\omega_n^{k-n/2}} p_1(\omega_{n/2}^{k-n/2}) & \text{if } k \geq n/2 \end{cases} \\
 \omega_n^k &= -\omega_n^{k-\frac{n}{2}}
 \end{aligned}$$

Thus, if $k < n/2$:

$$\begin{aligned}
 p_0(\omega_{n/2}^k) + \omega_n^k p_1(\omega_{n/2}^k) &= p(\omega_n^k) \\
 p_0(\omega_{n/2}^k) - \omega_n^k p_1(\omega_{n/2}^k) &= p(\omega_n^{k+n/2})
 \end{aligned}$$

A further improvement

Example:

$$p(\omega_4^0) = p_0(\omega_2^0) + \omega_4^0 p_1(\omega_2^0)$$

$$p(\omega_4^1) = p_0(\omega_2^1) + \omega_4^1 p_1(\omega_2^1)$$

$$p(\omega_4^2) = p_0(\omega_2^0) - \omega_4^0 p_1(\omega_2^0)$$

$$p(\omega_4^3) = p_0(\omega_2^1) - \omega_4^1 p_1(\omega_2^1)$$

6. Fast Fourier Transform

Algorithm: FFT

Input: Array a containing the n coefficients of a polynomial p and $n = 2^k$

Output: $DFT_n(p)$

FFT (a , n)

1. **if** $n = 1$ **then** /* p is constant */
2. **return** a $\swarrow p^0$
3. $d^{[0]} = FFT([a_0, a_2, \dots, a_{n-2}], n/2)$
4. $d^{[1]} = FFT([a_1, a_3, \dots, a_{n-1}], n/2)$
5. $\omega_n = e^{2\pi i/n} \quad \nwarrow p^1$
6. $\omega = 1$
7. **for** $k = 0$ **to** $n/2 - 1$ **do** { /* $\omega = \underline{\underline{\omega_n^k}}$ */
8. $d_k = \underline{d_k^{[0]}} + \omega \cdot \underline{d_k^{[1]}}$
9. $d_{k+n/2} = \underline{d_k^{[0]}} - \omega \cdot \underline{d_k^{[1]}}$
10. $\omega = \underline{\omega} \cdot \omega$
11. **return** d

FFT: Example

$$p(x) = 3x^3 - 15x^2 + 18x + 0$$

$$a = [0, 18, -15, 3]$$

$$a^{[0]} = [0, -15] \quad a^{[1]} = [18, 3]$$

$$\begin{aligned} FFT([0, -15], 2) &= (FFT([0], 1) + FFT([-15], 1), \quad FFT([0], 1) - FFT([-15], 1)) \\ &= (-15, 15) \end{aligned}$$

$$\begin{aligned} FFT([18, 3], 2) &= (FFT([18], 1) + FFT([3], 1), \quad FFT([18], 1) - FFT([3], 1)) \\ &= (21, 15) \end{aligned}$$

$$k = 0 ; \omega = 1$$

$$d_0 = -15 + 1 * 21 = 6 \quad d_2 = -15 - 1 * 21 = -36$$

$$k = 1 ; \omega = i$$

$$d_1 = 15 + i * 15 \quad d_3 = 15 - i * 15$$

$$FFT(a, 4) = (6, 15+15i, -36, 15-15i)$$

7. Analysis

$T(n)$ = Time required for evaluating a polynomial of degree $< n$ at the points $\omega_{2n}^0, \omega_{2n}^1, \dots, \omega_{2n}^{2n-1}$.

$$\begin{aligned}
 T(1) &= O(1) \\
 T(n) &= 2 \cdot T(n/2) + O(n) \\
 &= O(n \log n)
 \end{aligned}$$

Conquer
Divide & Merge

Polynomial multiplication

Compute the product of two polynomials p, q of degree $< n$:

p, q of degree $n-1$, n coefficients



Evaluation via FFT: $\omega_{2n}^0, \omega_{2n}^1, \dots, \omega_{2n}^{2n-1}$ $\mathcal{O}(n \log n)$

$2n$ point-value pairs $(\omega_{2n}^i, p(\omega_{2n}^i))$ and $(\omega_{2n}^i, q(\omega_{2n}^i))$



Pointwise multiplication $\mathcal{O}(n)$

$2n$ point-value pairs $(\omega_{2n}^i, pq(\omega_{2n}^i))$



Interpolation

{
}

pq of degree $2n-2$, $2n-1$ coefficients

Interpolation

Convert the point-value representation into coefficient representation.

Input: $(\underline{x_0}, y_0), \dots, (\underline{x_{n-1}}, y_{n-1})$ where $x_i \neq x_j$, for all $i \neq j$

Output: Polynomial p with coefficients $\underline{a_0, \dots, a_{n-1}}$,
such that

$$\begin{aligned}
 p(\underline{x_0}) &= a_0 + a_1 x_0 + \dots + a_{n-1} x_0^{n-1} = \underline{y_0} \\
 p(x_1) &= a_0 + a_1 x_1 + \dots + a_{n-1} x_1^{n-1} = y_1 \\
 p(x_2) &= a_0 + a_1 x_2 + \dots + a_{n-1} x_2^{n-1} = y_2 \\
 &\vdots \\
 p(x_{n-1}) &= a_0 + a_1 x_{n-1} + \dots + a_{n-1} x_{n-1}^{n-1} = y_{n-1}
 \end{aligned}$$

unknown *given*

Interpolation

Matrix notation:

$$\begin{array}{c} \nearrow \\ \nabla_n \end{array} \quad \left(\begin{array}{cccc} 1 & x_0 & x_0^2 & \cdots & x_0^{n-1} \\ 1 & x_1 & x_1^2 & \cdots & x_1^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_{n-1} & x_{n-1}^2 & \cdots & x_{n-1}^{n-1} \end{array} \right) \left(\begin{array}{c} a_0 \\ a_1 \\ \vdots \\ a_{n-1} \end{array} \right) = \left(\begin{array}{c} y_0 \\ y_1 \\ \vdots \\ y_{n-1} \end{array} \right)$$

a_i γ

$$a_0 + a_1 \cdot x_0 + a_2 \cdot x_0^2 + \dots + a_{n-1} \cdot x_0^{n-1} = \gamma_0$$

, ; ;

Interpolation

System of equations

$$\forall_n \begin{array}{c} \\ \downarrow \\ \end{array} \begin{array}{cccc} 1 & x_0 & \cdots & x_0^{n-1} \\ 1 & x_1 & \cdots & x_1^{n-1} \\ \vdots & x_{i-1}^i & \ddots & \vdots \\ 1 & x_{n-1} & \cdots & x_{n-1}^{n-1} \end{array} \left(\begin{array}{c} a_0 \\ a_1 \\ \vdots \\ a_{n-1} \end{array} \right) = \left(\begin{array}{c} y_0 \\ y_1 \\ \vdots \\ y_{n-1} \end{array} \right)$$

solvable if $x_i \neq x_j$ for all $i \neq j$.

Special case (here) : $x_i = \omega_n^i$ $x_i^j = (\omega_n^i)^j = \omega_n^{ij}$

Definition: $V_n = \underline{\underline{\omega_n^{ij}}}_{i,j}, \quad a = (a_i), \quad y = (y_i)$

$$V_n a = y \quad \Rightarrow \quad a = V_n^{-1} y$$

$$V_n^{-1} \cdot V_n a = V_n^{-1} y$$

Interpolation

Theorem:

For any $0 \leq i, j \leq n - 1$ we have:

$$(V_n^{-1})_{ij} = \frac{\omega_n^{-ij}}{n}$$

Proof:

$$V_n^{-1} = \underbrace{\left(\frac{\omega_n^{-ij}}{n} \right)_{i,j}}$$

We have to show:

$$\underbrace{V_n^{-1} V_n}_{\text{Identity}} = I_n^{\checkmark}$$

Interpolation

Consider the entry of $V_n^{-1}V_n$ in row i and column j :

$$\begin{aligned}
 & (V_n^{-1}V_n)_{ij} = \\
 & \left(\frac{1}{n}, \frac{\omega_n^{-i}}{n}, \dots, \frac{\omega_n^{-i(n-1)}}{n} \right) \begin{pmatrix} \dots & 1 & \dots \\ \dots & \omega_n^j & \dots \\ \dots & \omega_n^{2j} & \dots \\ \vdots & \vdots & \vdots \\ \dots & \omega_n^{(n-1)j} & \dots \end{pmatrix}_{ij} = \\
 & = \sum_{k=0}^{n-1} \frac{1}{n} \cdot \omega_n^{-ik} \cdot \omega_n^{jk} \\
 & = \frac{1}{n} \cdot \sum_{k=0}^{n-1} \omega_n^{(-i+j)k} \\
 & = \begin{cases} 1 & i = j \\ 0 & \text{otherwise} \end{cases}
 \end{aligned}$$

Interpolation

$$\left(V_n^{-1} V_n \right)_{ij} = \sum_{k=0}^{n-1} \frac{\omega_n^{-ik}}{n} \omega_n^{jk} = \frac{1}{n} \sum_{k=0}^{n-1} \omega_n^{(-i+j)k} = I_n$$

Case 1: $i = j$

$$\frac{1}{n} \sum_{k=0}^{n-1} \omega_n^{(-i+j)k} \stackrel{=\sigma}{=} \frac{1}{n} \sum_{k=0}^{n-1} \underbrace{\omega_n^{0 \cdot k}}_{=1} = 1 \quad \checkmark$$

Case 2: $i \neq j$, i.e. $-(n-1) \leq -i + j \leq n-1$
 $\underline{\hspace{1cm}}$ thus $n \nmid -i + j$:

$$\frac{1}{n} \sum_{k=0}^{n-1} \omega_n^{(-i+j)k} = 0 \quad \checkmark$$

Interpolation

Summation lemma:

For any integer $n > 0$, $l \geq 0$ with $\underline{n \nmid l}$:

$$\sum_{k=0}^{n-1} \omega_n^{lk} = 0$$

Geometric Series

$$\sum_{k=0}^{n-1} x^k = \frac{x^n - 1}{x - 1}$$

Proof:

$$\sum_{k=0}^{n-1} (\omega_n^l)^k = \frac{(\underbrace{\omega_n^l}_\text{! !})^n - 1}{\omega_n^l - 1} = \frac{(\omega_n^n)^l - 1}{\omega_n^l - 1} = 0$$

ω_n^0

Interpolation

$$\begin{aligned}
 a_i &= \left(V_n^{-1} y \right)_i \\
 &= \underbrace{\left(\frac{1}{n}, \frac{\omega_n^{-i}}{n}, \dots, \frac{\omega_n^{-i(n-1)}}{n} \right)}_{\longrightarrow} \begin{pmatrix} y_0 \\ y_1 \\ \vdots \\ y_{n-1} \end{pmatrix} \quad \downarrow \\
 &= \sum_{k=0}^{n-1} y_k \frac{\omega_n^{-ik}}{n}
 \end{aligned}$$

$$r(x) := \sum_{k=0}^{n-1} y_k \cdot x^k$$

$$= \frac{1}{n} \sum_{k=0}^{n-1} y_k \left(\omega_n^{-i} \right)^k = \frac{1}{n} \cdot r(\omega_n^{-i})$$

$\leadsto \text{DFT} !$

Interpolation

$$a = \frac{1}{n} \left(\underbrace{\sum_{k=0}^{n-1} y_k (\omega_n^{-0})^k}, \underbrace{\sum_{k=0}^{n-1} y_k (\omega_n^{-1})^k}, \dots, \underbrace{\sum_{k=0}^{n-1} y_k (\omega_n^{-(n-1)})^k} \right)$$

$$r(x) = y_0 + y_1 x + y_2 x^2 + \dots + y_{n-1} x^{n-1}$$

$$a = \frac{1}{n} (r(\omega_n^{-0}), r(\omega_n^{-1}), \dots, r(\omega_n^{-(n-1)}))$$

Interpolation and DFT

$$a = \frac{1}{n} (r(\omega_n^{-0}), r(\omega_n^{-1}), \dots, r(\omega_n^{-(n-1)}))$$

$$a = \frac{1}{n} (r(\omega_n^n), r(\omega_n^{n-1}), \dots, r(\omega_n^1)) \quad \text{since } \omega_n^n = 1$$

$$\rightarrow a_{\circled{i}} = \frac{1}{n} (DFT_n(r))_{\circled{n-i}} \quad (i \neq 0)$$

$$\rightarrow a_{\circled{0}} = \frac{1}{n} (DFT_n(r))_{\circled{0}}$$

Polynomial multiplication by FFT

Compute the product of two polynomials p, q of degree $< n$:

p, q of degree $n-1$, n coefficients

