

Theoretical Computer Science - Bridging Course

Summer Term 2018

Exercise Sheet 9

for getting feedback submit electronically by 06:00 am, Monday, July 2nd, 2018

Exercise 1: Propositional Logic: Basic Terms *(2+2+2+2 Points)*

Let $\Sigma := \{p, q, r\}$ be a set of atoms. An interpretation $I : \Sigma \rightarrow \{T, F\}$ maps every atom to either true or false. Inductively, an interpretation I can be extended to composite formulae φ over Σ (cf. lecture). We write $I \models \varphi$ if φ evaluates to T (true) under I . In case $I \models \varphi$, I is called a *model* for φ .

For each of the following formulae, give *all* interpretations which are models. Make a truth table and/or use logical equivalencies to find all models (document your steps). Which of these formulae are satisfiable, which are unsatisfiable and which are tautologies?

(a) $\varphi_1 = (p \wedge \neg q) \vee (\neg p \vee q)$

(b) $\varphi_2 = (\neg p \wedge (\neg p \vee q)) \leftrightarrow (p \vee \neg q)$

(c) $\varphi_3 = (p \wedge \neg q) \rightarrow \neg(p \wedge q)$

(d) $\varphi_4 = (p \wedge q) \rightarrow (p \vee r)$

Remark: $a \rightarrow b := \neg a \vee b$, $a \leftrightarrow b := (a \rightarrow b) \wedge (b \rightarrow a)$, $a \not\rightarrow b := \neg(a \rightarrow b)$.

Sample Solution

(a) See Table 1. The result shows that φ_1 is a tautology.

(b) See Table 2. The result shows that φ_2 is satisfiable.

(c) φ_3 is equivalent to $\neg(p \wedge \neg q) \vee (\neg p \vee \neg q)$ which is equivalent to $(\neg p \vee q) \vee (\neg p \vee \neg q)$ which is equivalent to $\neg p \vee q \vee \neg p \vee \neg q$ which is equivalent to $\neg p \vee \neg q \vee q$ which is a tautology as either q or $\neg q$ holds.

(d) See Table 3. The result shows that φ_4 is tautology.

model	p	q	$p \wedge \neg q$	$\neg p \vee q$	φ_1
✓	0	0	0	1	1
✓	0	1	0	1	1
✓	1	0	1	1	1
✗	1	1	0	1	1

Table 1: Truthtables for Exercises 1 (a).

model	p	q	$\neg p \vee q$	$\neg p \wedge (\neg p \vee q)$	$p \vee \neg q$	φ_2
✓	0	0	1	1	1	1
✗	0	1	1	1	0	0
✗	1	0	0	0	1	0
✗	1	1	1	0	1	0

Table 2: Truthtables for Exercises 1 (b).

model	p	q	r	$p \wedge q$	$p \vee r$	φ_4
✓	0	0	0	0	0	1
✓	0	0	1	0	1	1
✓	0	1	0	0	0	1
✓	0	1	1	0	1	1
✓	1	0	0	0	1	1
✓	1	0	1	0	1	1
✓	1	1	0	1	1	1
✓	1	1	1	1	1	1

Table 3: Truthtables for Exercises 1 (d).

Exercise 2: CNF and DNF

(2+2 Points)

- (a) Convert $\varphi_1 := (p \rightarrow q) \rightarrow (\neg r \wedge q)$ into Conjunctive Normal Form (CNF).
(b) Convert $\varphi_2 := \neg((\neg p \rightarrow \neg q) \wedge \neg r)$ into Disjunctive Normal Form (DNF).

Remark: Use the known logical equivalencies given in the lecture slides to do the necessary transformations. State which equivalency you are using in each step.

Sample Solution

(a)

$$\begin{aligned} & (p \rightarrow q) \rightarrow (\neg r \wedge q) \\ \equiv & \neg(\neg p \vee q) \vee (\neg r \wedge q) && \text{Definition of '}\rightarrow\text{'} \\ \equiv & (p \wedge \neg q) \vee (\neg r \wedge q) && \text{De Morgan} \\ \equiv & ((p \wedge \neg q) \vee \neg r) \wedge ((p \wedge \neg q) \vee q) && \text{Distribution} \\ \equiv & ((p \vee \neg r) \wedge (\neg q \vee \neg r)) \wedge ((p \vee q) \wedge (\neg q \vee q)) && \text{Distribution} \\ \equiv & ((p \vee \neg r) \wedge (\neg q \vee \neg r)) \wedge ((p \vee q) \wedge 1) && \text{Complementation} \\ \equiv & ((p \vee \neg r) \wedge (\neg q \vee \neg r)) \wedge (p \vee q) && \text{Identity} \\ \equiv & (p \vee \neg r) \wedge (\neg q \vee \neg r) \wedge (p \vee q) && \text{Associativity} \end{aligned}$$

(b)

$$\begin{aligned} & \neg((\neg p \rightarrow \neg q) \wedge \neg r) \\ \equiv & \neg((p \vee \neg q) \wedge \neg r) && \text{Definition of '}\rightarrow\text{'} \\ \equiv & \neg(p \vee \neg q) \vee r && \text{De Morgan} \\ \equiv & (\neg p \wedge q) \vee r && \text{De Morgan} \end{aligned}$$

Exercise 3: Logical Entailment

(2+2 Points)

A *knowledge base* KB is a set of formulae over a given set of atoms Σ . An interpretation I of Σ is called a *model* of KB , if it is a model for *all* formulae in KB . A knowledge base KB *entails* a formula φ (we write $KB \models \varphi$), if *all* models of KB are also models of φ .

Let $KB := \{p \vee q, \neg r \vee p\}$. Show or disprove that KB logically entails the following formulae.

- (a) $\varphi_1 := (p \wedge q) \vee \neg(\neg r \vee p)$
(b) $\varphi_2 := (q \leftrightarrow r) \rightarrow p$

Sample Solution

- (a) KB does not entail φ_1 . Consider the interpretation $I : p \mapsto 1, q \mapsto 0, r \mapsto 0$. Interpretation I is a model for KB but not for φ_1 .
(b) Table 4 shows that every model of KB is also a model of φ_2 , hence $KB \models \varphi_2$.

model of KB	p	q	r	$p \vee q$	$\neg r \vee p$	$q \leftrightarrow r$	φ_2	model of φ_2
\times	0	0	0	0	0	1	0	\times
\times	0	0	1	0	0	0	1	\checkmark
\checkmark	0	1	0	1	1	0	1	\checkmark
\times	0	1	1	1	0	1	0	\times
\checkmark	1	0	0	1	1	1	1	\checkmark
\checkmark	1	0	1	1	1	0	1	\checkmark
\checkmark	1	1	0	1	1	0	1	\checkmark
\checkmark	1	1	1	1	1	1	1	\checkmark

Table 4: Truthtable for Exercise 3 (b).

Exercise 4: Inference Rules and Calculi

(2+2 Points)

Let $\varphi_1, \dots, \varphi_n, \psi$ be propositional formulae. An *inference rule*

$$\frac{\varphi_1, \dots, \varphi_n}{\psi}$$

means that if $\varphi_1, \dots, \varphi_n$ are 'considered true', then ψ is 'considered true' as well ($n = 0$ is the special case of an axiom). A (propositional) *calculus* \mathbf{C} is described by a *set* of inference rules.

Given a formula ψ and knowledge base $KB := \{\varphi_1, \dots, \varphi_n\}$ (where $\varphi_1, \dots, \varphi_n$ are formulae) we write $KB \vdash_{\mathbf{C}} \psi$ if ψ can be derived from KB by starting from a subset of KB and repeatedly applying inference rules from the calculus \mathbf{C} to 'generate' new formulae until ψ is obtained.

Consider the following two calculi, defined by their inference rules (φ, ψ, χ are arbitrary formulae).

$$\mathbf{C}_1 : \frac{\varphi \rightarrow \psi, \psi \rightarrow \chi}{\varphi \rightarrow \chi}, \frac{\neg\varphi \rightarrow \psi}{\neg\psi \rightarrow \varphi}, \frac{\varphi \leftrightarrow \psi}{\varphi \rightarrow \psi, \psi \rightarrow \varphi}$$

$$\mathbf{C}_2 : \frac{\varphi, \varphi \rightarrow \psi}{\psi}, \frac{\varphi \wedge \psi}{\varphi, \psi}, \frac{(\varphi \wedge \psi) \rightarrow \chi}{\varphi \rightarrow (\psi \rightarrow \chi)}$$

Using the respective calculus, show the following derivations (document your steps).

(a) $\{p \leftrightarrow \neg r, \neg q \rightarrow r\} \vdash_{\mathbf{C}_1} p \rightarrow q$

(b) $\{p \wedge q, p \rightarrow r, (q \wedge r) \rightarrow s\} \vdash_{\mathbf{C}_2} s$

Remark: Inferences of a given calculus are purely syntactical, i.e. rules only apply in their specific form (much like a grammar) and no other logical transformations not given in the calculus are allowed.

Sample Solution

(a) We use \mathbf{C}_1 to derive new formulae until we obtain the desired one.

$$\begin{array}{l} \neg q \rightarrow r \quad \text{2nd rule} \\ \vdash_{\mathbf{C}_1} \quad \neg r \rightarrow q \\ \\ p \leftrightarrow \neg r \quad \text{3rd rule} \\ \vdash_{\mathbf{C}_1} \quad p \rightarrow \neg r, \neg r \rightarrow p \\ \\ p \rightarrow \neg r, \neg r \rightarrow q \quad \text{1st rule} \\ \vdash_{\mathbf{C}_1} \quad p \rightarrow q \end{array}$$

(b) We use \mathbf{C}_2 to derive new formulae until we obtain the desired one.

$$\begin{array}{rcl}
 p \wedge q & \begin{array}{c} \text{2nd rule} \\ \vdash_{\mathbf{C}_2} \end{array} & p, q \\
 p, p \rightarrow r & \begin{array}{c} \text{1st rule} \\ \vdash_{\mathbf{C}_2} \end{array} & r \\
 (q \wedge r) \rightarrow s & \begin{array}{c} \text{3rd rule} \\ \vdash_{\mathbf{C}_2} \end{array} & q \rightarrow (r \rightarrow s) \\
 q, q \rightarrow (r \rightarrow s) & \begin{array}{c} \text{1st rule} \\ \vdash_{\mathbf{C}_2} \end{array} & r \rightarrow s \\
 r, r \rightarrow s & \begin{array}{c} \text{1st rule} \\ \vdash_{\mathbf{C}_2} \end{array} & s
 \end{array}$$