




Chapter 6

Graph Algorithms

Algorithm Theory
WS 2015/16

Fabian Kuhn

Flow Network



Ford Fulkerson: Running Time

- Time of regular Ford-Fulkerson algorithm with integer capacities:

$$O(mC)$$

- Time of algorithm with scaling parameter:

$$O(m^2 \log C)$$

- $O(\log C)$ is polynomial in the size of the input, but not in n
- Can we get an algorithm that runs in time polynomial in n ?
- Always picking a **shortest augmenting path** leads to running time

$$O(m^2 n)$$

Other Maximum Flow Algorithms

- There are many other algorithms to solve the maximum flow problem, for example:
- **Preflow-push algorithm:**
 - Maintains a preflow (\forall nodes: inflow \geq outflow)
 - Alg. guarantees: As soon as we have a flow, it is optimal
 - Detailed discussion in 2012/13 lecture
 - Running time of basic algorithm: $O(m \cdot n^2)$
 - Doing steps in the “right” order: $O(n^3)$
- **Current best known complexity: $O(m \cdot n)$**
 - For graphs with $m \geq n^{1+\epsilon}$ [King,Rao,Tarjan 1992/1994]
(for every constant $\epsilon > 0$)
 - For sparse graphs with $m \leq n^{16/15-\delta}$ [Orlin, 2013]

Maximum Flow Applications

- Maximum flow has many applications
- Reducing a problem to a max flow problem can even be seen as an important algorithmic technique
- Examples:
 - related network flow problems
 - computation of small cuts
 - computation of matchings
 - computing disjoint paths
 - scheduling problems
 - assignment problems with some side constraints
 - ...

Undirected Edges and Vertex Capacities

Undirected Edges:


- Undirected edge $\{u, v\}$: add edges (u, v) and (v, u) to network

Vertex Capacities:

- Not only edges, but also (or only) nodes have capacities
- Capacity c_v of node $v \notin \{s, t\}$:

$$f^{\text{in}}(v) = f^{\text{out}}(v) \leq c_v$$

- Replace node v by edge $e_v = \{v_{\text{in}}, v_{\text{out}}\}$:



Minimum s - t Cut

Given: undirected graph $G = (V, E)$, nodes $s, t \in V$

s - t cut: Partition (A, B) of V such that $s \in A, t \in B$

Size of cut (A, B) : number of edges crossing the cut

Objective: find s - t cut of minimum size

Edge Connectivity

Definition: A graph $G = (V, E)$ is **k -edge connected** for an integer $k \geq 1$ if the graph **$G_X = (V, E \setminus X)$ is connected** for every edge set

$$X \subseteq E, |X| \leq k - 1.$$

Goal: Compute **edge connectivity $\lambda(G)$** of G
(and edge set X of size $\lambda(G)$ that divides G into ≥ 2 parts)

- minimum set X is a minimum s - t cut for some $s, t \in V$
 - Actually for all s, t in different components of $G_X = (V, E \setminus X)$
- Possible algorithm: fix s and find min s - t cut for all $t \neq s$

Minimum s - t Vertex-Cut

Given: undirected graph $G = (V, E)$, nodes $s, t \in V$

s - t vertex cut: Set $X \subset V$ such that $s, t \notin X$ and s and t are in different components of the sub-graph $G[V \setminus X]$ induced by $V \setminus X$

Size of vertex cut: $|X|$

Objective: find s - t vertex-cut of minimum size

- Replace undirected edge $\{u, v\}$ by (u, v) and (v, u)
- Compute max s - t flow for edge capacities ∞ and node capacities

$$c_v = 1 \text{ for } v \neq s, t$$

- Replace each node v by v_{in} and v_{out} :
- Min edge cut corresponds to min vertex cut in G

Vertex Connectivity

Definition: A graph $G = (V, E)$ is **k -vertex connected** for an integer $k \geq 1$ if the sub-graph $G[V \setminus X]$ **induced by $V \setminus X$ is connected** for every edge set

$$X \subseteq V, |X| \leq k - 1.$$

Goal: Compute **vertex connectivity $\kappa(G)$** of G
(and node set X of size $\kappa(G)$ that divides G into ≥ 2 parts)

- Compute minimum s - t vertex cut for fixed s and all $t \neq s$

Edge-Disjoint Paths

Given: Graph $G = (V, E)$ with nodes $s, t \in V$

Goal: Find as many edge-disjoint s - t paths as possible

Solution:

- Find max s - t flow in G with **edge capacities** $c_e = 1$ for all $e \in E$

Flow f induces **$|f|$ edge-disjoint paths:**

- Integral capacities \rightarrow can compute integral max flow f
- Get $|f|$ edge-disjoint paths by greedily picking them
- Correctness follows from flow conservation $f^{\text{in}}(v) = f^{\text{out}}(v)$

Vertex-Disjoint Paths

Given: Graph $G = (V, E)$ with nodes $s, t \in V$

Goal: Find as many internally vertex-disjoint s - t paths as possible

Solution:

- Find max s - t flow in G with **node capacities** $c_v = 1$ for all $v \in V$

Flow f induces **$|f|$ vertex-disjoint paths:**

- Integral capacities \rightarrow can compute integral max flow f
- Get $|f|$ vertex-disjoint paths by greedily picking them
- Correctness follows from flow conservation $f^{\text{in}}(v) = f^{\text{out}}(v)$

Menger's Theorem

Theorem: (edge version)

For every graph $G = (V, E)$ with nodes $s, t \in V$, the size of the minimum s - t (edge) cut equals the maximum number of pairwise edge-disjoint paths from s to t .

Theorem: (node version)

For every graph $G = (V, E)$ with nodes $s, t \in V$, the size of the minimum s - t vertex cut equals the maximum number of pairwise internally vertex-disjoint paths from s to t .

- Both versions can be seen as a special case of the max flow min cut theorem