



Chapter 6

Graph Algorithms

Algorithm Theory
WS 2018/19

Fabian Kuhn

Circulations with Demands

Given: Directed network with positive edge capacities

Sources & Sinks: Instead of one source and one destination, several sources that generate flow and several sinks that absorb flow.

Supply & Demand: sources have supply values, sinks demand values

Goal: Compute a flow such that source supplies and sink demands are exactly satisfied

- The circulation problem is a feasibility rather than a maximization problem

Circulations with Demands: Formally

Given: Directed network $G = (V, E)$ with

- Edge capacities $c_e > 0$ for all $e \in E$
- Node demands $\underline{d_v} \in \mathbb{R}$ for all $v \in V$
 - $\underline{d_v} > 0$: node needs flow and therefore is a sink
 - $d_v < 0$: node has a supply of $-d_v$ and is therefore a source
 - $d_v = 0$: node is neither a source nor a sink


Flow: Function $f: E \rightarrow \mathbb{R}_{\geq 0}$ satisfying

- *Capacity Conditions:* $\forall e \in E: 0 \leq f(e) \leq c_e$
- *Demand Conditions:* $\forall v \in V: \underline{f^{\text{in}}(v) - f^{\text{out}}(v) = d_v}$


Objective: Does a flow f satisfying all conditions exist?
If yes, find such a flow f .

Reduction to Maximum Flow

- Add “super-source” s^* and “super-sink” t^* to network



Example



Circulation with Demands


Theorem: There is a feasible circulation with demands $d_v, v \in V$ on graph G if and only if there is a flow of value D on G' .

- If all capacities and demands are integers, there is an integer circulation

The **max flow min cut theorem** also implies the following:

Theorem: The graph G has a feasible circulation with demands $d_v, v \in V$ if and only if for all cuts (A, B) ,

$$\sum_{v \in B} d_v \leq c(A, B).$$



Circulation: Demands and Lower Bounds

Given: Directed network $G = (V, E)$ with


- Edge capacities $c_e > 0$ and **lower bounds** $0 \leq \underline{\ell}_e \leq \underline{c}_e$ for $e \in E$
- Node demands $d_v \in \mathbb{R}$ for all $v \in V$
 - $d_v > 0$: node needs flow and therefore is a sink
 - $d_v < 0$: node has a supply of $-d_v$ and is therefore a source
 - $d_v = 0$: node is neither a source nor a sink

Flow: Function $f: E \rightarrow \mathbb{R}_{\geq 0}$ satisfying

- *Capacity Conditions:* $\forall e \in E: \ell_e \leq f(e) \leq c_e$
- *Demand Conditions:* $\forall v \in V: f^{\text{in}}(v) - f^{\text{out}}(v) = d_v$

Objective: Does a flow f satisfying all conditions exist?
If yes, find such a flow f .

Solution Idea



- Define **initial circulation** $f_0(e) = \underline{\ell}_e$
 Satisfies capacity constraints: $\forall e \in E: \underline{\ell}_e \leq f_0(e) \leq \underline{c}_e$

$$L_v = 12 - 7 = 5$$

- Define

$$\underline{L}_v := \underline{f_0^{\text{in}}(v)} - \underline{f_0^{\text{out}}(v)} = \sum_{e \text{ into } v} \ell_e - \sum_{e \text{ out of } v} \ell_e$$


- If $\underline{L}_v = \underline{d}_v$, demand condition is satisfied at v by f_0 , otherwise, we need to superimpose another circulation f_1 such that

$$\underline{d}'_v := \underline{f_1^{\text{in}}(v)} - \underline{f_1^{\text{out}}(v)} = \underline{d}_v - \underline{L}_v$$

- Remaining capacity of edge e : $c'_e := c_e - \ell_e$
 $0 \leq f_1(e) \leq c'_e$
 $f_0(e) = \ell_e$
 $f_1(e) = f_0(e) + f_1(e) \leq c_e$
- We get a circulation problem with new demands \underline{d}'_v , new capacities \underline{c}'_e , and **no lower bounds**

Eliminating a Lower Bound: Example

Lower bound of 2




Reduce to Problem Without Lower Bounds

Graph $G = (V, E)$:

- Capacity: For each edge $e \in E: \ell_e \leq f(e) \leq c_e$
- Demand: For each node $v \in V: f^{\text{in}}(v) - f^{\text{out}}(v) = d_v$

Model lower bounds with supplies & demands:



Create Network G' (without lower bounds):

- For each edge $e \in E: \underline{c'_e} = \underline{c_e - \ell_e}$
- For each node $v \in V: \underline{d'_v} = \underline{d_v - \underline{L_v}}$

Circulation: Demands and Lower Bounds

Theorem: There is a feasible circulation in \underline{G} (with lower bounds) if and only if there is feasible circulation in \underline{G}' (without lower bounds).

- Given circulation f' in G' , $f(e) = f'(e) + \ell_e$ is circulation in G
 - The capacity constraints are satisfied because $f'(e) \leq c_e - \ell_e$
 - Demand conditions:

$$\begin{aligned} \underline{f^{\text{in}}(v) - f^{\text{out}}(v)} &= \sum_{e \text{ into } v} (\ell_e + f'(e)) - \sum_{e \text{ out of } v} (\ell_e + f'(e)) \\ &= \underline{L_v} + (d_v - L_v) = d_v \end{aligned}$$

- Given circulation f in G , $f'(e) = f(e) - \ell_e$ is circulation in G'
 - The capacity constraints are satisfied because $\ell_e \leq f(e) \leq c_e$
 - Demand conditions:

$$\begin{aligned} \underline{f'^{\text{in}}(v) - f'^{\text{out}}(v)} &= \sum_{e \text{ into } v} (f(e) - \ell_e) - \sum_{e \text{ out of } v} (f(e) - \ell_e) \\ &= \underline{d_v - L_v} \end{aligned}$$

Integrality

Theorem: Consider a circulation problem with integral capacities, flow lower bounds, and node demands. If the problem is feasible, then it also has an integral solution.

Proof:

- Graph G' has only integral capacities and demands
- Thus, the flow network used in the reduction to solve circulation with demands and no lower bounds has only integral capacities
- The theorem now follows because a max flow problem with integral capacities also has an optimal integral solution
- It also follows that with the max flow algorithms we studied, we get an integral feasible circulation solution.


Matrix Rounding

- **Given:** $p \times q$ matrix $D = \{d_{i,j}\}$ of real numbers
- **row i sum:** $a_i = \sum_j d_{i,j}$, **column j sum:** $b_j = \sum_i d_{i,j}$
- **Goal:** **Round** each $d_{i,j}$, as well as a_i and b_j up or down to the next integer so that the sum of rounded elements in each row (column) equals the rounded row (column) sum
- **Original application:** publishing census data

Example:

<u>3.14</u>	<u>6.80</u>	<u>7.30</u>	<u>17.24</u>
9.60	2.40	0.70	<u>12.70</u>
3.60	1.20	6.50	<u>11.30</u>
<u>16.34</u>	<u>10.40</u>	<u>14.50</u>	

original data



<u>3</u>	<u>7</u>	<u>7</u>	<u>17</u>
10	2	1	<u>13</u>
3	1	7	<u>11</u>
<u>16</u>	<u>10</u>	<u>15</u>	

possible rounding

Matrix Rounding

Theorem: For any matrix, there exists a feasible rounding.

Remark: Just rounding to the nearest integer doesn't work

<u>0.35</u>	<u>0.35</u>	<u>0.35</u>	1.05
0.55	0.55	<u>0.55</u>	1.65
0.90	0.90	0.90	

original data

0	0	0	0
<u>1</u>	<u>1</u>	<u>1</u>	3
<u>1</u>	<u>1</u>	<u>1</u>	

rounding to nearest integer


0	0	1	1
1	1	0	2
1	1	1	

feasible rounding

Reduction to Circulation

<u>3.14</u>	<u>6.80</u>	<u>7.30</u>	<u>17.24</u>
9.60	2.40	0.70	12.70
3.60	1.20	6.50	11.30
16.34	<u>10.40</u>	14.50	

Matrix elements and row/column sums give a feasible circulation that satisfies all lower bound, capacity, and demand constraints



Matrix Rounding

Theorem: For any matrix, there exists a feasible rounding.


Proof:

- The matrix entries $d_{i,j}$ and the row and column sums a_i and b_j give a feasible circulation for the constructed network
- Every feasible circulation gives matrix entries with corresponding row and column sums (follows from demand constraints)
- Because all demands, capacities, and flow lower bounds are integral, there is an integral solution to the circulation problem

→ gives a feasible rounding!


Gifts-Children Graph

- Which child likes which gift can be represented by a graph




Matching

Matching: Set of pairwise non-incident edges



Maximal Matching: A matching s.t. no more edges can be added

Maximum Matching: A matching of maximum possible size




Perfect Matching: Matching of size $\frac{n}{2}$ (every node is matched)

Bipartite Graph

Definition: A graph $G = (V, E)$ is called bipartite iff its node set can be partitioned into two parts $V = V_1 \cup V_2$ such that for each edge $\{u, v\} \in E$,

$$|\{u, v\} \cap V_1| = 1.$$

- Thus, edges are only between the two parts




Santa's Problem

Maximum Matching in Bipartite Graphs:

Every child can get a gift
iff there is a matching
of size $\#$ children


Clearly, every matching
is at most as big

If $\#$ children = $\#$ gifts,
there is a solution iff
there is a perfect matching



Reducing to Maximum Flow

- Like edge-disjoint paths...



all capacities are 1

Reducing to Maximum Flow

Theorem: Every integer solution to the max flow problem on the constructed graph induces a maximum bipartite matching of G .

Proof:

1. An integer flow f of value $|f|$ induces a matching of size $|f|$
 - Left nodes (gifts) have incoming capacity 1
 - Right nodes (children) have outgoing capacity 1
 - Left and right nodes are incident to ≤ 1 edge e of G with $f(e) = 1$
2. A matching of size k implies a flow f of value $|f| = k$
 - For each edge $\{u, v\}$ of the matching:
$$f((s, u)) = f((u, v)) = f((v, t)) = 1$$
 - All other flow values are 0


Running Time of Max. Bipartite Matching

Theorem: A maximum matching of a bipartite graph can be computed in time $O(\underline{m \cdot n})$.


Use Ford-Fulkerson
time: $O(m \cdot C)$
 $C \leq n/2$
value of max. flow

Perfect Matching?

- There can only be a perfect matching if both sides of the partition have size $n/2$.
- There is no perfect matching, iff there is an $s-t$ cut of size $< n/2$ in the flow network.



s - t Cuts



Partition (A , B) of node set such that $s \in A$ and $t \in B$

- If $v_i \in \underline{A}$: edge (v_i, t) is in cut (A, B)
- If $u_i \in B$: edge (s, u_i) is in cut (A, B)
- Otherwise (if $u_i \in A$, $v_i \in B$), all edges from u_i to some $v_j \in B$ are in cut (A, B)

Hall's Marriage Theorem


Theorem: A bipartite graph $G = (U \cup V, E)$ for which $|U| = |V|$ has a perfect matching if and only if

$$\forall U' \subseteq U: |N(U')| \geq |U'|,$$

where $N(U') \subseteq V$ is the set of neighbors of nodes in U' .

Proof: No perfect matching \Leftrightarrow some s - t cut has capacity $< n/2$

1. Assume there is U' for which $|N(U')| < |U'|$:



Hall's Marriage Theorem


Theorem: A bipartite graph $G = (U \cup V, E)$ for which $|U| = |V|$ has a perfect matching if and only if

$$\forall U' \subseteq U: |N(U')| \geq |U'|,$$

where $N(U') \subseteq V$ is the set of neighbors of nodes in U' .

Proof: No perfect matching \Leftrightarrow some s - t cut has capacity $< n/2$

2. Assume that there is a cut (A, B) of capacity $< n/2$



Hall's Marriage Theorem

Theorem: A bipartite graph $G = (U \cup V, E)$ for which $|U| = |V|$ has a perfect matching if and only if

$$\forall U' \subseteq U: |N(U')| \geq |U'|,$$

where $N(U') \subseteq V$ is the set of neighbors of nodes in U' .

Proof: No perfect matching \Leftrightarrow some s - t cut has capacity $< n$


2. Assume that there is a cut (A, B) of capacity $< n$

$$\left. \begin{array}{l} |U'| = \frac{n}{2} - x \\ |N(U')| \leq y + z \\ \hline x + y + z < \frac{n}{2} \end{array} \right\} \begin{array}{l} \frac{n}{2} - x > y + z \geq |N(U')| \\ |U'| \\ \hline \underline{|N(U')|} < \underline{|U'|} \end{array}$$

What About General Graphs

- Can we efficiently compute a maximum matching if G is not bipartite?
- How good is a **maximal matching**?
 - A matching that cannot be extended...
- **Vertex Cover**: set $S \subseteq V$ of nodes such that

$$\forall \{u, v\} \in E, \quad \{u, v\} \cap S \neq \emptyset.$$



- A vertex cover covers all edges by incident nodes


Vertex Cover vs Matching

Consider a matching M and a vertex cover S

Claim: $|M| \leq |S|$

Proof:

- At least one node of every edge $\{u, v\} \in M$ is in S
- Needs to be a different node for different edges from M





Vertex Cover vs Matching

Consider a matching M and a vertex cover S $|M| \leq |S|$

Claim: If M is maximal and S is minimum, $|S| \leq 2|M|$

Proof:

- M is maximal: for every edge $\{u, v\} \in E$, either u or v (or both) are matched



- Every edge $e \in E$ is “covered” by at least one matching edge
- Thus, the set of the nodes of all matching edges gives a vertex cover S of size $|S| = 2|M|$.

Maximal Matching Approximation

Theorem: For any maximal matching M and any maximum matching M^* , it holds that

$$\underline{|M|} \geq \frac{|M^*|}{2}$$

Proof:

S^* : minimum vertex cover

$$|M^*| \leq |S^*| \leq 2|M|$$

Theorem: The set of all matched nodes of a maximal matching M is a vertex cover of size at most twice the size of a min. vertex cover.