



# Algorithms and Data Structures

## Winter Term 2020/2021

### Sample Solution Exercise Sheet 5

#### Exercise 1: Bad Hash Functions

Let  $m$  be the size of a hash table and  $M \gg m$  the largest possible key of the elements we want to store in the table. The following “hash functions” are poorly chosen. Explain for each function why it is not a suitable hash function.

- (a)  $h : x \mapsto \lfloor \frac{x}{m} \rfloor \bmod m$
- (b)  $h : x \mapsto (2x + 1) \bmod m$  ( $m$  even).
- (c)  $h : x \mapsto (x \bmod m) + \lfloor \frac{m}{x+1} \rfloor$
- (d) For each calculation of the hash value of  $x$  one chooses for  $h(x)$  a uniform random number from  $\{0, \dots, m-1\}$
- (e)  $h : x \mapsto \lfloor \frac{M}{x \cdot p \bmod M} \rfloor \bmod m$ , where  $p$  is prime and  $\frac{M}{2} < p < M$
- (f) For a set of “good” hash functions  $h_1, \dots, h_\ell$  with  $\ell \in \Theta(\log m)$ , we first compute  $h_1(x)$ , then  $h_2(h_1(x))$  etc. until  $h_\ell(h_{\ell-1}(\dots h_1(x)) \dots)$ . That is, the function is  $h : k \mapsto h_\ell(h_{\ell-1}(\dots h_1(x)) \dots)$

#### Sample Solution

- (a) Values are not scattered.  $m$  subsequent values have the same hash value.
- (b) Only half of the hash table is used. The cells  $0, 2, 4, \dots, m-2$  stay empty.
- (c)  $h(m-1) = m$ , but the table has only the positions  $0, \dots, m-1$ .
- (d) The hash value of  $x$  can not be reproduced.
- (e) First, consider the function  $h' : x \mapsto \lfloor \frac{M}{x} \rfloor \bmod m$ .  $h'$  maps all  $x > M/2$  (i.e., half of the keys) to position 1, all  $x$  with  $M/3 < x \leq M/2$  (i.e. 1/6 of the keys) to position 2 etc. So the table is filled asymmetrically. As the function  $x \mapsto x \cdot p \bmod M$  is a bijection from  $\{0, \dots, M-1\}$  to  $\{0, \dots, M-1\}$ ,  $h$  has the same property of an asymmetrical filled table (but compared to  $h'$  we do not have that a long sequence of subsequent keys are mapped to the same position which would be another undesirable property, cf. part (a)).
- (f) The calculation of a single hash value needs  $\Omega(\log m)$ .

## Exercise 2: (No) Families of Universal Hash Functions

- (a) Let  $\mathcal{S} = \{0, \dots, M-1\}$  and  $\mathcal{H}_1 := \{h : x \mapsto a \cdot x^2 \pmod{m} \mid a \in \mathcal{S}\}$ . Show that  $\mathcal{H}_1$  is not  $c$ -universal for constant  $c \geq 1$  (that is  $c$  is fixed and must not depend on  $m$ ).
- (b) Let  $m$  be a prime and let  $k = \lfloor \log_m M \rfloor$ . We consider the keys  $x \in \mathcal{S}$  in base  $m$  presentation, i.e.,  $x = \sum_{i=0}^k x_i m^i$ . Consider the set of functions from the lecture (week 5, slide 15)

$$\mathcal{H}_2 := \left\{ h : x \mapsto \sum_{i=0}^k a_i x_i \pmod{m} \mid a_i \in \{0, \dots, m-1\} \right\}.$$

Show that  $\mathcal{H}_2$  is 1-universal.

*Hint: Two keys  $x \neq y$  have to differ at some digit  $x_j \neq y_j$  in their base  $m$  representation.*

## Sample Solution

- (a) For an  $x \in \mathcal{S}$  let  $y = x + i \cdot m \in \mathcal{S}$  for some  $i \in \mathbb{Z} \setminus \{0\}$ . Such a  $y$  exists for any  $x$  if  $M > 2m$ . Let  $h \in \mathcal{H}_1$ . We obtain

$$\begin{aligned} h(y) &= a \cdot y^2 \pmod{m} \\ &\equiv a \cdot (x + im)^2 \pmod{m} \\ &\equiv a \cdot (x^2 + 2xim + (im)^2) \pmod{m} \\ &\equiv ax^2 \pmod{m} = h(x). \end{aligned}$$

It follows that  $|\{h \in \mathcal{H}_1 \mid h(x) = h(y)\}| = |\mathcal{H}_1|$ , so for  $m > c$  we have

$$|\{h \in \mathcal{H}_1 \mid h(x) = h(y)\}| > \frac{c}{m} |\mathcal{H}_1|.$$

This means that for  $m > c$ ,  $\mathcal{H}_1$  is not  $c$ -universal.

- (b) Let  $x, y \in \mathcal{S}$  with  $x \neq y$ . Let  $x_j \neq y_j$  be the position where  $x$  and  $y$  differ in their base  $m$  representation. Let  $h \in \mathcal{H}_2$  such that  $h(x) = h(y)$ . We have

$$\begin{aligned} h(x) &= h(y) \\ \iff \sum_{i=0}^k a_i x_i &\equiv \sum_{i=0}^k a_i y_i \pmod{m} \\ \iff a_j \underbrace{(x_j - y_j)}_{\neq 0} &\equiv \sum_{i \neq j} a_i (y_i - x_i) \pmod{m} \\ \iff a_j &\equiv (x_j - y_j)^{-1} \sum_{i \neq j} a_i (y_i - x_i) \pmod{m} \quad (x_j - y_j)^{-1} \text{ exists because } m \text{ is prime} \end{aligned}$$

This means that for any values  $a_0, \dots, a_{j-1}, a_{j+1}, \dots, a_k$  there is a unique  $a_j$  such that the function  $h$  defined by  $a_0, \dots, a_k$  is in  $\{h \in \mathcal{H}_2 \mid h(x) = h(y)\}$ . So we have  $m^k$  possibilities to choose a function from  $\{h \in \mathcal{H}_2 \mid h(x) = h(y)\}$ . It follows

$$\frac{|\{h \in \mathcal{H}_2 \mid h(x) = h(y)\}|}{|\mathcal{H}_2|} = \frac{m^k}{m^{k+1}} = \frac{1}{m}.$$